

WO04014426

Publication Title:

WO04014426

Abstract:

Abstract not available for WO04014426 Data supplied from the esp@cenet database - Worldwide

Courtesy of <http://v3.espacenet.com>

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
19 February 2004 (19.02.2004)

PCT

(10) International Publication Number
WO 2004/014426 A1

- (51) International Patent Classification⁷: **A61K 45/00**, 31/517, A61P 35/00, 9/00
- (74) Agent: **ASTRAZENECA**; Global Intellectual Property, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).
- (21) International Application Number: **PCT/GB2003/003390**
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: **5 August 2003 (05.08.2003)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data: **0218526.2** **9 August 2002 (09.08.2002)** **GB**
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except MG, US*): **ASTRAZENECA AB [SE/SE]; Sodertalje, S-151 85 (SE).**
- (71) Applicant (*for MG only*): **ASTRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).**
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): **WEDGE, Stephen, Robert [GB/GB]; AstraZeneca R & D Alderley, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).**
- Published:
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **COMBINATION OF VEGF RECEPTOR TYROSINE KINASE INHIBITORS FOR TREATMENT OF CANCER**

(57) Abstract: The present invention relates to a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation, particularly a method for the treatment of a cancer, particularly a cancer involving a solid tumor, which comprises the administration of ZD6474 in combination with ZD1839; to a pharmaceutical composition comprising ZD6474 and ZD1839; to a combination product comprising ZD6474 and ZD1839 for use in a method of treatment of a human or animal body by therapy; to a kit comprising ZD6474 and ZD1839; to the use of ZD6474 and ZD1839 in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation.

WO 2004/014426 A1

COMBINATION OF VEGF RECEPTOR TYROSINE KINASE INHIBITORS FOR TREATMENT OF
CANCER


The present invention relates to a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation, particularly a method for the treatment of a cancer, particularly a cancer involving a solid tumour, which comprises the administration of ZD6474 in combination with ZD1839; to a pharmaceutical composition comprising ZD6474 and ZD1839; to a combination product comprising ZD6474 and ZD1839 for use in a method of treatment of a human or animal body by therapy; to a kit comprising ZD6474 and ZD1839; to the use of ZD6474 and ZD1839 in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is optionally being treated with ionising radiation.

Normal angiogenesis plays an important role in a variety of processes including embryonic development, wound healing and several components of female reproductive function. Undesirable or pathological angiogenesis has been associated with disease states including diabetic retinopathy, psoriasis, cancer, rheumatoid arthritis, atheroma, Kaposi's sarcoma and haemangioma (Fan et al, 1995, Trends Pharmacol. Sci. 16: 57-66; Folkman, 1995, Nature Medicine 1: 27-31). Alteration of vascular permeability is thought to play a role in both normal and pathological physiological processes (Cullinan-Bove et al, 1993, Endocrinology 133: 829-837; Senger et al, 1993, Cancer and Metastasis Reviews, 12: 303-324). Several polypeptides with *in vitro* endothelial cell growth promoting activity have been identified including, acidic and basic fibroblast growth factors (aFGF & bFGF) and vascular endothelial growth factor (VEGF). By virtue of the restricted expression of its receptors, the growth factor activity of VEGF, in contrast to that of the FGFs, is relatively specific towards endothelial cells. Recent evidence indicates that VEGF is an important stimulator of both normal and pathological angiogenesis (Jakeman et al, 1993, Endocrinology, 133: 848-859; Kolch et al, 1995, Breast Cancer Research and Treatment, 36:139-155) and vascular permeability (Connolly et al, 1989, J. Biol. Chem. 264: 20017-20024). Antagonism of VEGF action by sequestration of VEGF with antibody can result in inhibition of tumour growth (Kim et al, 1993, Nature 362: 841-844).

Receptor tyrosine kinases (RTKs) are important in the transmission of biochemical signals across the plasma membrane of cells. These transmembrane molecules characteristically consist of an extracellular ligand-binding domain connected through a segment in the plasma membrane to an intracellular tyrosine kinase domain. Binding of ligand to the receptor results in

stimulation of the receptor-associated tyrosine kinase activity which leads to phosphorylation of tyrosine residues on both the receptor and other intracellular molecules. These changes in tyrosine phosphorylation initiate a signalling cascade leading to a variety of cellular responses. To date, at least nineteen distinct RTK subfamilies, defined by amino acid sequence homology, have been identified. One of these subfamilies is presently comprised by the *fms*-like tyrosine kinase receptor, Flt-1, the kinase insert domain-containing receptor, KDR (also referred to as Flk-1), and another *fms*-like tyrosine kinase receptor, Flt-4. Two of these related RTKs, Flt-1 and KDR, have been shown to bind VEGF with high affinity (De Vries et al, 1992, Science 255: 989-991; Terman et al, 1992, Biochem. Biophys. Res. Comm. 1992, 187: 1579-1586). Binding of VEGF to these receptors expressed in heterologous cells has been associated with changes in the tyrosine phosphorylation status of cellular proteins and calcium fluxes.

Quinazoline derivatives which are inhibitors of VEGF receptor tyrosine kinase are described in International Patent Applications Publication Nos. WO 98/13354 and WO 01/32651. In WO 98/13354 and WO 01/32651 compounds are described which possess activity against VEGF receptor tyrosine kinase (VEGF RTK) whilst possessing some activity against epidermal growth factor (EGF) receptor tyrosine kinase (EGF RTK). ZD6474 is 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline:



ZD6474

ZD6474 falls within the broad general disclosure of WO 98/13354 and is exemplified in WO 01/32651. ZD6474 is a potent inhibitor of VEGF RTK and also has some activity against EGF RTK. ZD6474 has been shown to elicit broad-spectrum anti-tumour activity in a range of models following once-daily oral administration (Wedge S.R., Ogilvie D.J., Dukes M. et al, Proc. Am. Assoc. Canc. Res. 2001; 42: abstract 3126).

In WO 98/13354 and WO 01/32651 it is stated that compounds of their inventions:
“may be applied as a sole therapy or may involve, in addition to a compound of the invention,
one or more other substances and/or treatments. Such conjoint treatment may be achieved by
way of the simultaneous, sequential or separate administration of the individual components of
5 the treatment.”

WO 98/13354 and WO 01/32651 then go on to describe examples of such conjoint treatment
including surgery, radiotherapy and various types of chemotherapeutic agent including inhibitors
of growth factor function.


Nowhere in WO 98/13354 and WO 01/32651 does it suggest the combination of a
10 compound of the invention and an epidermal growth factor receptor tyrosine kinase inhibitor
for the treatment of any disease state including cancer.

Nowhere in WO 98/13354 and WO 01/32651 is the specific combination of ZD6474
and ZD1839 suggested.

Nowhere in WO 98/13354 and WO 01/32651 does it state that use of any compound
15 of the invention therein with other treatments will produce surprisingly beneficial effects.

Unexpectedly and surprisingly we have now found that the particular compound
ZD6474 used in combination with a particular selection from the broad description of
combination therapies listed in WO 98/13354 and WO 01/32651, namely with ZD1839,
produces significantly better effects than any one of ZD6474 and ZD1839 used alone. In
20 particular, ZD6474 used in combination with ZD1839 produces significantly better effects on
solid tumours than any one of ZD6474 and ZD1839 used alone.

ZD1839 is *N*-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine:



25

ZD1839

ZD1839 is also known as gefitinib and is also known as Iressa™ (Trademark of
AstraZeneca UK Limited) and it is an epidermal growth factor receptor (EGFR) tyrosine

kinase inhibitor (TKI). ZD1839 is described in International Patent Application Publication No. WO 96/33980.

In recent years it has been discovered that certain growth factor tyrosine kinase enzymes are important in the transmission of biochemical signals which initiate cell replication.

5 They are large proteins which span the cell membrane and possess an extracellular binding domain for growth factors, for example the epidermal growth factor receptor (EGFR) which binds epidermal growth factor (EGF), and an intracellular portion which functions as a kinase to phosphorylate tyrosine amino acids in proteins and hence to influence cell proliferation.

EGFR is a member of the erbB family of receptor tyrosine kinases, which includes

10 EGFR, erbB2, erbB3 and erbB4, and it is known that these receptor tyrosine kinases are frequently involved in driving the proliferation and survival of tumour cells (reviewed in Olayioye et al., EMBO J., 2000, 19, 3159). One mechanism by which this can be accomplished is by overexpression of the receptor at the protein level, generally as a result of gene amplification. This has been observed in many common human cancers (reviewed in

15 Klapper et al., Adv. Cancer Res., 2000, 77, 25), such as breast cancer (Sainsbury et al., Brit. J. Cancer, 1988, 58, 458; Guerin et al., Oncogene Res., 1988, 3, 21; Slamon et al., Science, 1989, 244, 707; Klijn et al., Breast Cancer Res. Treat., 1994, 29, 73 and reviewed in Salomon et al., Crit. Rev. Oncol. Hematol., 1995, 19, 183), non-small cell lung cancers (NSCLCs) including adenocarcinomas (Cerny et al., Brit. J. Cancer, 1986, 54, 265; Reubi et al., Int. J.

20 Cancer, 1990, 45, 269; Rusch et al., Cancer Research, 1993, 53, 2379; Brabender et al., Clin. Cancer Res., 2001, 7, 1850) as well as other cancers of the lung (Hendler et al., Cancer Cells, 1989, 7, 347; Ohsaki et al., Oncol. Rep., 2000, 7, 603), bladder cancer (Neal et al., Lancet, 1985, 366; Chow et al., Clin. Cancer Res., 2001, 7, 1957, Zhau et al., Mol Carcinog., 3, 254), oesophageal cancer (Mukaida et al., Cancer, 1991, 68, 142), gastrointestinal cancer such as

25 colon, rectal or stomach cancer (Bolen et al., Oncogene Res., 1987, 1, 149; Kapitanovic et al., Gastroenterology, 2000, 112, 1103; Ross et al., Cancer Invest., 2001, 19, 554), cancer of the prostate (Visakorpi et al., Histochem. J., 1992, 24, 481; Kumar et al., 2000, 32, 73; Scher et al., J. Natl. Cancer Inst., 2000, 92, 1866), leukaemia (Konaka et al., Cell, 1984, 37, 1035, Martin-Subero et al., Cancer Genet Cytogenet., 2001, 127, 174), ovarian cancer (Hellstrom et

30 al., Cancer Res., 2001, 61, 2420), head and neck cancer (Shiga et al., Head Neck, 2000, 22, 599) and pancreatic cancer (Ovotny et al., Neoplasma, 2001, 48, 188).

It is widely believed that as a consequence of the dysfunctional regulation of one or more of these receptors many tumours become clinically more aggressive and this correlates with a poorer prognosis for the patient (Brabender et al, Clin. Cancer Res., 2001, 7, 1850; Ross et al, Cancer Investigation, 2001, 19, 554, Yu et al., Bioessays, 2000, 22.7, 673). In addition to these clinical findings, a wealth of pre-clinical information suggests that the erbB family of receptor tyrosine kinases are involved in cellular transformation. This includes the observations that many tumour cell lines overexpress one or more of the erbB receptors and that EGFR or erbB2 when transfected into non-tumour cells have the ability to transform these cells. In addition to this, a number of pre-clinical studies have demonstrated that anti-proliferative effects can be induced by knocking out one or more erbB activities by small molecule inhibitors, dominant negatives or inhibitory antibodies (reviewed in Mendelsohn et al., Oncogene, 2000, 19, 6550). Thus it has been recognised that inhibitors of these receptor tyrosine kinases should be of value as selective inhibitors of the proliferation of mammalian cancer cells (Yaish *et al.* Science, 1988, 242, 933, Kolibaba *et al.*, Biochimica et Biophysica Acta, 1997, 133, F217-F248; Al-Obeidi *et al.*, 2000, Oncogene, 19, 5690-5701; Mendelsohn *et al.*, 2000, Oncogene, 19, 6550-6565). In addition to this pre-clinical data, the use of inhibitory antibodies against EGFR and erbB2 (c-225 and trastuzumab respectively) has proven to be beneficial in the clinic for the treatment of selected solid tumours (reviewed in Mendelsohn *et al.*, 2000, Oncogene, 19, 6550-6565).

It is believed that members of the erbB type receptor tyrosine kinase family may be implicated in a number of non-malignant proliferative disorders because amplification and/or activity of erbB receptor tyrosine kinases has been detected in psoriasis (Ben-Bassat, Curr. Pharm. Des., 2000, 6, 933; Elder et al., Science, 1989, 243, 811), benign prostatic hyperplasia (BPH) (Kumar et al., Int. Urol. Nephrol., 2000, 32,73), atherosclerosis and restenosis (Bokemeyer et al., Kidney Int., 2000, 58, 549). It is therefore expected that inhibitors of erbB type receptor tyrosine kinases will be useful in the treatment of these and other non-malignant disorders involving excessive cellular proliferation.

It is known from International Patent Application Publication No. WO 96/33980 that ZD1839 possesses EGF RTK inhibitory activity (J R Woodburn *et al.* in Proc. Amer. Assoc. Canc. Res., 1997, 38, 633 and Pharmacol. Ther., 1999, 82, 241-250) and is an inhibitor of the proliferation of cancer tissue.

It is stated in WO 96/33980 that compounds of the invention, which include ZD1839, may be given conjointly with other cancer therapies. It states therein

"The anti-proliferative treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the quinazoline derivative of the invention, one or more other

- 5 anti-tumour substances, for example cytotoxic or cytostatic anti-tumour substances, for example those selected from, for example, mitotic inhibitors, for example vinblastine, vindesine and vinorelbine; tubulin disassembly inhibitors such as taxol; alkylating agents, for example cis-platin, carboplatin and cyclophosphamide; antimetabolites, for example 5-fluorouracil, tegafur, methotrexate, cytosine arabinoside and hydroxyurea, or, for example, one of the
- 10 preferred antimetabolites disclosed in European Patent Application No. 239362 such as N-{5-[N-(3,4-dihydro-2-methyl-4-oxoquinazolin-6-ylmethyl)-N-methylamino]-2-thenoyl}-L-glutamic acid; intercalating antibiotics, for example adriamycin, mitomycin and bleomycin; enzymes, for example asparaginase; topoisomerase inhibitors, for example etoposide and camptothecin; biological response modifiers, for example interferon;
- 15 anti-hormones, for example antioestrogens such as tamoxifen, for example antiandrogens such as 4'-cyano-3-(4-fluorophenylsulphonyl)-2-hydroxy-2-methyl-3'-(trifluoromethyl)-propionanilide or, for example LHRH antagonists or LHRH agonists such as goserelin, leuporelin or buserelin and hormone synthesis inhibitors, for example aromatase inhibitors such as those disclosed in European Patent Application No. 0296749, for example 2,2'-[5-(1H-
- 20 1,2,4- triazol-1-ylmethyl)-1,3-phenylene]bis(2-methylpropionitrile), and, for example, inhibitors of 5 α -reductase such as 17 β -(N-tert-butylcarbonyl)-4-aza-5 α -androst-1-en-3-one."

Nowhere in WO 96/33980 does it suggest any combination of an EGF RTK inhibitor with a VEGF RTK inhibitor for the treatment of any disease state including cancer.

- International Patent Application Publication No. WO 02/41882 describes, generally,
- 25 combinations of agents which decrease VEGF activity and agents which decrease EGF activity. A study has been carried out examining co-administration of antibodies to KDR and EGFR (Shaheen, R.M., et al., Brit. J. Cancer, 2001, 85, 584-589).

- Unexpectedly and surprisingly we have now found that the particular compound ZD6474 used in combination with the particular compound ZD1839, produces significantly
- 30 better anti-tumour effects than each of ZD6474 and ZD1839 used alone.

Anti-cancer effects of a method of treatment of the present invention include, but are not limited to, anti-tumour effects, the response rate, the time to disease progression and the

survival rate. Anti-tumour effects of a method of treatment of the present invention include but are not limited to, inhibition of tumour growth, tumour growth delay, regression of tumour, shrinkage of tumour, increased time to regrowth of tumour on cessation of treatment, slowing of disease progression. It is expected that when a method of treatment of the present
5 invention is administered to a warm-blooded animal such as a human, in need of treatment for cancer, with or without a solid tumour, said method of treatment will produce an effect, as measured by, for example, one or more of: the extent of the anti-tumour effect, the response rate, the time to disease progression and the survival rate.

According to the present invention there is provided a method for the production of an
10 antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof.

According to a further aspect of the present invention there is provided a method for
15 the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof.

According to a further aspect of the present invention there is provided a method for
20 the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof.

According to a further aspect of the present invention there is provided a method for
25 the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof; wherein ZD6474 and ZD1839 may each optionally be administered together with a
30 pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises

administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof; wherein ZD6474 and ZD1839 may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

- 5 According to a further aspect of the present invention there is provided a method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof; wherein ZD6474 and
- 10 ZD1839 may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

- According to a further aspect of the invention there is provided a pharmaceutical composition which comprises ZD6474 or a pharmaceutically acceptable salt thereof, and ZD1839 or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically
- 15 acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a combination product comprising ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof, for use in a method of treatment of a human or animal body by therapy.

- 20 According to a further aspect of the present invention there is provided a kit comprising ZD6474 or a pharmaceutically acceptable salt thereof, and ZD1839 or a pharmaceutically acceptable salt thereof.

According to a further aspect of the present invention there is provided a kit comprising:

- 25 a) ZD6474 or a pharmaceutically acceptable salt thereof in a first unit dosage form;
b) ZD1839 or a pharmaceutically acceptable salt thereof in a second unit dosage form;
and
c) container means for containing said first and second dosage forms.

- According to a further aspect of the present invention there is provided a kit
- 30 comprising:

- a) ZD6474 or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable excipient or carrier, in a first unit dosage form;

- b) ZD1839 or a pharmaceutically acceptable salt thereof together with a pharmaceutically acceptable excipient or carrier, in a second unit dosage form; and
c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided the use of
5 ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human.

According to a further aspect of the present invention there is provided the use of
10 ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as a human.

According to a further aspect of the present invention there is provided the use of
ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically
15 acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-tumour effect in a warm-blooded animal such as a human.

According to a further aspect of the present invention there is provided a therapeutic combination treatment comprising the administration of an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, optionally together with a pharmaceutically
20 acceptable excipient or carrier, and the simultaneous, sequential or separate administration of an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof; wherein ZD1839 may optionally be administered together with a pharmaceutically acceptable excipient or carrier, to a warm-blooded animal such as a human in need of such therapeutic treatment.

Such therapeutic treatment includes an antiangiogenic and/or vascular permeability
25 effect, an anti-cancer effect and an anti-tumour effect.

A combination treatment of the present invention as defined herein may be achieved by way of the simultaneous, sequential or separate administration of the individual components of said treatment. A combination treatment as defined herein may be applied as a sole therapy or may involve surgery or radiotherapy or an additional chemotherapeutic agent, in addition to a
30 combination treatment of the invention.

Surgery may comprise the step of partial or complete tumour resection, prior to, during or after the administration of the combination treatment with ZD6474 described herein.

Other chemotherapeutic agents for optional use with a combination treatment of the present invention include those described in WO 01/32651 which is incorporated herein by reference. Such chemotherapy may cover five main categories of therapeutic agent:

- (i) other antiangiogenic agents including vascular targeting agents;
- 5 (ii) cytostatic agents;
- (iii) biological response modifiers (for example interferon);
- (iv) antibodies (for example edrecolomab); and
- (v) antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology.

10 The administration of a triple combination of ZD6474, ZD1839 and ionising radiation may produce effects, such as anti-tumour effects, greater than those achieved with any of ZD6474, ZD1839 and ionising radiation used alone, greater than those achieved with the combination of ZD6474 and ZD1839, greater than those achieved with the combination of ZD6474 and ionising radiation and greater than those achieved with the combination of
15 ZD1839 and ionising radiation.

According to the present invention there is provided a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective
20 amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable
25 salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human,
30 which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective

amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation.

According to a further aspect of the present invention there is provided a method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation, wherein ZD6474 and ZD1839 may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation, wherein ZD6474 and ZD1839 may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided a method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation, wherein ZD6474 and ZD1839 may each optionally be administered together with a pharmaceutically acceptable excipient or carrier.

According to a further aspect of the present invention there is provided the use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

According to a further aspect of the present invention there is provided the use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically

acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

According to a further aspect of the present invention there is provided the use of
5 ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-tumour effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

According to a further aspect of the present invention there is provided a therapeutic
10 combination treatment comprising the administration of an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, optionally together with a pharmaceutically acceptable excipient or carrier, and the administration of an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof, optionally together with a pharmaceutically acceptable excipient or carrier and the administration of an effective amount of ionising
15 radiation, to a warm-blooded animal such as a human in need of such therapeutic treatment wherein the ZD6474, ZD1839 and ionising radiation may be administered simultaneously, sequentially or separately and in any order.

A warm-blooded animal such as a human which is being treated with ionising radiation means a warm-blooded animal such as a human which is treated with ionising radiation before,
20 after or at the same time as the administration of a medicament or combination treatment comprising ZD6474 and ZD1839. For example said ionising radiation may be given to said warm-blooded animal such as a human within the period of a week before to a week after the administration of a medicament or combination treatment comprising ZD6474 and ZD1839. This means that ZD6474, ZD1839 and ionising radiation may be administered separately or
25 sequentially in any order, or may be administered simultaneously. The warm-blooded animal may experience the effect of each of ZD6474, ZD1839 and radiation simultaneously.

According to one aspect of the present invention the ionising radiation is administered before one of ZD6474 and ZD1839 or after one of ZD6474 and ZD1839.

According to one aspect of the present invention the ionising radiation is administered
30 before both ZD6474 and ZD1839 or after both ZD6474 and ZD1839.

According to one aspect of the present invention ZD6474 is administered to a warm-blooded animal after the animal has been treated with ionising radiation.

In another aspect of the present invention ZD6474 and ZD1839 are dosed daily continuously for a longer period of time during which time ionising radiation is administered periodically, that is for a few days, for example 1, 2, 3, 4 or 5 days at a time.

According to another aspect of the present invention the effect of a method of
5 treatment of the present invention is expected to be at least equivalent to the addition of the effects of each of the components of said treatment used alone, that is, of each of ZD6474 and ZD1839 or of each of ZD6474, ZD1839 and ionising radiation used alone.

According to another aspect of the present invention the effect of a method of
treatment of the present invention is expected to be greater than the addition of the effects of
10 each of the components of said treatment used alone, that is, of each of ZD6474 and ZD1839 or of each of ZD6474, ZD1839 and ionising radiation, used alone.

According to another aspect of the present invention the effect of a method of
treatment of the present invention is expected to be a synergistic effect.

It should also be appreciated that according to the present invention a combination
15 treatment is defined as affording a synergistic effect if the effect is therapeutically superior, as measured by, for example, the extent of the response, the response rate, the time to disease progression or the survival period, to that achievable on dosing one or other of the components of the combination treatment at its conventional dose. For example, the effect of the combination treatment is synergistic if the effect is therapeutically superior to the effect
20 achievable with ZD6474 or ZD1839 or ionising radiation alone. Further, the effect of the combination treatment is synergistic if a beneficial effect is obtained in a group of patients that does not respond (or responds poorly) to ZD6474 or ZD1839 or ionising radiation alone. In addition, the effect of the combination treatment is defined as affording a synergistic effect if one of the components is dosed at its conventional dose and the other component(s) is/are
25 dosed at a reduced dose and the therapeutic effect, as measured by, for example, the extent of the response, the response rate, the time to disease progression or the survival period, is equivalent to that achievable on dosing conventional amounts of the components of the combination treatment. In particular, synergy is deemed to be present if the conventional dose of ZD6474 or ZD1839 or ionising radiation may be reduced without detriment to one or more
30 of the extent of the response, the response rate, the time to disease progression and survival data, in particular without detriment to the duration of the response, but with fewer and/or less

troublesome side-effects than those that occur when conventional doses of each component are used.

As stated above the combination treatments of the present invention as defined herein are of interest for their antiangiogenic and/or vascular permeability effects. Combination
5 treatments of the invention are expected to be useful in the prophylaxis and treatment of a wide range of disease states where inappropriate angiogenesis occurs including cancer, diabetes, psoriasis, rheumatoid arthritis, Kaposi's sarcoma, haemangioma, acute and chronic nephropathies, atheroma, arterial restenosis, autoimmune diseases, acute inflammation, lymphoedema, endometriosis, dysfunctional uterine bleeding and ocular diseases with retinal
10 vessel proliferation including macular degeneration. In particular such combination treatments of the invention are expected to slow advantageously the growth of primary and recurrent solid tumours of, for example, the colon, breast, prostate, lungs and skin. More especially combination treatments of the present invention are expected to slow advantageously the growth of tumours in lung cancer, particularly non-small cell lung cancer (NSCLC). More
15 particularly such combination treatments of the invention are expected to inhibit any form of cancer associated with VEGF including leukaemia, multiple myeloma and lymphoma and also, for example, to inhibit the growth of those primary and recurrent solid tumours which are associated with VEGF, especially those tumours which are significantly dependent on VEGF for their growth and spread, including for example, certain tumours of the colon, breast,
20 prostate, lung, vulva and skin, particularly NSCLC.

In another aspect of the present invention ZD6474 and ZD1839, optionally with ionising radiation, are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with EGF especially those tumours which are significantly dependent on EGF for their growth and spread.

25 In another aspect of the present invention ZD6474 and ZD1839, optionally with ionising radiation, are expected to inhibit the growth of those primary and recurrent solid tumours which are associated with both VEGF and EGF especially those tumours which are significantly dependent on VEGF and EGF for their growth and spread.

The compositions described herein may be in a form suitable for oral administration, for
30 example as a tablet or capsule, for nasal administration or administration by inhalation, for example as a powder or solution, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) for example as a sterile solution, suspension or

emulsion, for topical administration for example as an ointment or cream, for rectal administration for example as a suppository or the route of administration may be by direct injection into the tumour or by regional delivery or by local delivery. In other embodiments of the present invention the ZD6474 of the combination treatment may be delivered

5 endoscopically, intratracheally, intralesionally, percutaneously, intravenously, subcutaneously, intraperitoneally or intratumourally. Preferably ZD6474 is administered orally. In general the compositions described herein may be prepared in a conventional manner using conventional excipients. The compositions of the present invention are advantageously presented in unit dosage form.

10 ZD6474 will normally be administered to a warm-blooded animal at a unit dose within the range 10-500mg per square metre body area of the animal, for example approximately 0.3-15mg/kg in a human. A unit dose in the range, for example, 0.3-15mg/kg, preferably 0.5-5mg/kg is envisaged and this is normally a therapeutically-effective dose. A unit dosage form such as a tablet or capsule will usually contain, for example 25-500mg of active
15 ingredient. Preferably a daily dose in the range of 0.5-5mg/kg is employed.

For ZD1839, a conventional tablet formulation may be used for oral administration to humans containing 50 mg, 100 mg, 250 mg or 500 mg of active ingredient. Conveniently the daily oral dose of ZD1839 is, for example, in the range 25 to 750 mg, preferably in the range 50 to 600 mg, more preferably in the range 100 to 400mg.

20 Radiotherapy may be administered according to the known practices in clinical radiotherapy. The dosages of ionising radiation will be those known for use in clinical radiotherapy. The radiation therapy used will include for example the use of γ -rays, X-rays, and/or the directed delivery of radiation from radioisotopes. Other forms of DNA damaging factors are also included in the present invention such as microwaves and UV-irradiation. For
25 example X-rays may be dosed in daily doses of 1.8-2.0Gy, 5 days a week for 5-6 weeks. Normally a total fractionated dose will lie in the range 45-60Gy. Single larger doses, for example 5-10Gy may be administered as part of a course of radiotherapy. Single doses may be administered intraoperatively. Hyperfractionated radiotherapy may be used whereby small doses of X-rays are administered regularly over a period of time, for example 0.1Gy per hour
30 over a number of days. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and on the uptake by cells.

As stated above the size of the dose of each therapy which is required for the therapeutic or prophylactic treatment of a particular disease state will necessarily be varied depending on the host treated, the route of administration and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is
5 treating any particular patient. For example, it may be necessary or desirable to reduce the above-mentioned doses of the components of the combination treatments in order to reduce toxicity.

The present invention relates to combinations of ZD1839 or a salt thereof with ZD6474 or with a salt of ZD6474.

10 Salts of ZD1839 for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of ZD1839 and its pharmaceutically acceptable salts. Salts include, for example, an acid-addition salt of ZD1839, for example, a mono- or di-acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric, maleic,
15 tartaric, fumaric, methanesulphonic or 4-toluenesulphonic acid.

Salts of ZD6474 for use in pharmaceutical compositions will be pharmaceutically acceptable salts, but other salts may be useful in the production of ZD6474 and its pharmaceutically acceptable salts. Such salts may be formed with an inorganic or organic base which affords a pharmaceutically acceptable cation. Such salts with inorganic or organic bases
20 include for example an alkali metal salt, such as a sodium or potassium salt, an alkaline earth metal salt such as a calcium or magnesium salt, an ammonium salt or for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

ZD6474 may be synthesised according to any of the known processes for making
25 ZD6474. For example ZD6474 may be made according to any of the processes described in WO 01/32651; for example those described in Examples 2(a), 2(b) and 2(c) of WO 01/32651.

ZD1839 may be synthesised according to any of the known processes for making ZD1839. For example ZD1839 may be made according to the processes described in WO 96/33980 (See Examples 1, 10 and 24-31).

30 The following tests were used to demonstrate the activity of ZD6474 in combination with ZD1839.

Human LoVo colon tumour xenografts in *Nude* mice

10⁷ LoVo tumour cells in 0.1 ml of serum free Dulbecco's Modified Eagle's Medium (DMEM) were injected subcutaneously (s.c.) into the flank of each athymic (*nu/nu* genotype) mouse. Tumor volumes were assessed by bilateral Vernier caliper measurement and, taking
5 length to be the longest diameter across the tumor and width the corresponding perpendicular, calculated using the formula (length x width) x the square root of (length x width) x ($\pi/6$). Five days after implantation when tumours reached a mean volume of approximately 0.3 cm³, mice were treated with ZD6474 (6 mg/kg/administration), ZD1839 (50 mg/kg/administration), or a combination thereof, orally (p.o.) daily for 15 days (day 0 – 14). ZD6474, ZD1839 or the
10 combination, were dosed at 0.1ml/10g body weight, as suspensions in 1% polysorbate 80 (i.e. a 1% (v/v) solution of polyoxyethylene (20) sorbitan mono-oleate in deionised water). Growth inhibition from the start of treatment was assessed by comparison of the differences in tumor volume between control and treated groups. A one-tailed two-sample *t*-test was used to evaluate the significance of tumour growth inhibition.

15

Table I

Treatment	% Inhibition of Tumour Growth at Day 14
ZD6474 (6 mg/kg/day, days 0 – 14, p.o.)	38 %
ZD1839 (50 mg/kg/day, days 0 – 14, p.o.)	50 %
ZD6474 + ZD1839	71 %

The combination of ZD6474 with ZD1839 produced greater inhibition of tumour growth than each of ZD6474 and ZD1839 alone.

20 The data is shown graphically in Figure 1.

Human A431 vulval carcinoma tumour xenografts in *Nude* mice

5 x 10⁶ A431 tumour cells in 0.1 ml of serum free Dulbecco's Modified Eagle's Medium (DMEM) were injected subcutaneously (s.c.) into the flank of each athymic (*nu/nu*

genotype) mouse. Eleven days after inoculation tumours were excised and dissected to generate cubic tumour fragments of $0.5 - 1 \text{ mm}^3$, which were implanted into further nude mice for a therapy experiment. Tumour volumes were assessed by bilateral Vernier caliper measurement and, taking length to be the longest diameter across the tumour and width the corresponding perpendicular, calculated using the formula $(\text{length} \times \text{width}) \times \text{the square root of } (\text{length} \times \text{width}) \times (\pi/6)$. Twenty eight days after implantation when tumours reached a mean volume of 0.7 cm^3 , mice were treated with ZD6474 (50 mg/kg/day), ZD1839 (50 mg/kg/day), or a combination thereof, orally (p.o.) for 31 days (day 28 – 59). ZD6474, ZD1839 or the combination, were dosed at 0.1ml/10g body weight, as suspensions in 1% polysorbate 80 (i.e. a 1% (v/v) solution of polyoxyethylene (20) sorbitan mono-oleate in deionised water). Inhibition of tumour growth from the start of treatment was assessed on day 49 (the point at which control animals were removed from the experiment because of tumour burden) by comparison of the differences in tumor volume between control and treated groups. In addition, the number of tumour regressions following 31 days of treatment was ascertained (tumour regression being evident if the tumour volume at day 59 was smaller than the pre-treatment value on day 28).

Table I

Treatment	Mean % Inhibition of tumour growth following 21 days of treatment (treatment from day 28 to day 49) [P value by two-tailed t-test]	Number of tumours regressing after 31 days of treatment (treatment from day 28 to day 59)
ZD6474 (50 mg/kg/day, days 28 – 59, p.o.)	116 % ($P < 0.001$) (i.e. 16% tumour regression)	6/9
ZD1839 (50 mg/kg/day, days 28 – 59, p.o.)	93% ($P = 0.001$)	3/9
ZD6474 + ZD1839	157% ($P < 0.001$) (i.e. 57% tumour regression)	9/9

The combination of ZD6474 with ZD1839 produced a greater inhibition of tumour growth than each of ZD6474 and ZD1839 alone at day 59 ($P < 0.001$ versus ZD6474 alone, and

P<0.009 versus ZD1839 alone; one-tailed t-test). In contrast to treatment with ZD6474 or ZD1839 alone, regression was induced in all A431 vulval carcinoma xenografts treated with the combination of ZD6474 and ZD1839. The magnitude of tumour regression induced by the combination (calculated by comparing the pre-treatment tumour volume (day 28) with the
5 volume following 31 days of treatment (day 59)), reached 71 ± 3 % (mean \pm SE) by the end of the experiment.

The data is shown graphically in Figure 2.

CLAIMS

1. A method for the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human, which comprises administering to
5 said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof.
2. A method for the production of an antiangiogenic and/or vascular permeability
10 reducing effect in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or simultaneously with an effective amount of ionising radiation.
15
3. A method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof.
20
4. A method for the treatment of a cancer in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or
25 simultaneously with an effective amount of ionising radiation.
5. A method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an
30 effective amount of ZD1839 or a pharmaceutically acceptable salt thereof.

6. A method for the treatment of a cancer involving a solid tumour in a warm-blooded animal such as a human, which comprises administering to said animal an effective amount of ZD6474 or a pharmaceutically acceptable salt thereof, before, after or simultaneously with an effective amount of ZD1839 or a pharmaceutically acceptable salt thereof and before, after or
5 simultaneously with an effective amount of ionising radiation.

7. A pharmaceutical composition which comprises ZD6474 or a pharmaceutically acceptable salt thereof, and ZD1839 or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable excipient or carrier.

10

8. A kit comprising ZD6474 or a pharmaceutically acceptable salt thereof, and ZD1839 or a pharmaceutically acceptable salt thereof.

9. Use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a
15 pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human.


10. Use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a
20 pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in a warm-blooded animal such as a human which is being treated with ionising radiation.


11. Use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a
25 pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-cancer effect in a warm-blooded animal such as a human.

12. Use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the
30 production of an anti-cancer effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

13. Use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-tumour effect in a warm-blooded animal such as a human.

- 5 14. Use of ZD6474 or a pharmaceutically acceptable salt thereof and ZD1839 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for use in the production of an anti-tumour effect in a warm-blooded animal such as a human which is being treated with ionising radiation.

**Figure 1**

**Figure 2**

INTERNATIONAL SEARCH REPORT

PCT/GB 03/03390

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K45/00 A61K31/517 A61P35/00 A61P9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

BIOSIS, WPI Data, EPO-Internal, PAJ, MEDLINE, EMBASE, COMPENDEX, SCISEARCH, CHEM
ABS Data, PASCAL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>MAGNÉ N ET AL: "Sequence-dependent effects of ZD1839 ('Iressa') in combination with cytotoxic treatment in human head and neck cancer" BRITISH JOURNAL OF CANCER 04 MAR 2002 UNITED KINGDOM, vol. 86, no. 5, 4 March 2002 (2002-03-04), pages 819-827, XP002256122 ISSN: 0007-0920 abstract; figures 3,5; table 4 page 821, column 1, last paragraph -column 2, paragraph 2 page 825, column 1, last line -page 826, column 1, paragraph 1 --- -/-</p>	1-14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

30 September 2003

Date of mailing of the international search report

16/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Laffargue-Haak, T

INTERNATIONAL SEARCH REPORT

PCT/GB 03/03390

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>CIARDIELLO F ET AL: "ANTITUMOR EFFECT AND POTENTIATION OF CYTOTOXIC DRUGS ACTIVITY IN HUMAN CANCER CELLS BY ZD-1839 (IRESSA), AN EPIDERMAL GROWTH FACTOR RECEPTOR-SELECTIVE TYROSINE KINASE INHIBITOR"</p> <p>CLINICAL CANCER RESEARCH, THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, US, vol. 6, no. 5, May 2000 (2000-05), pages 2053-2063, XP001009786</p> <p>ISSN: 1078-0432</p> <p>abstract</p> <p>page 2055, column 2, last paragraph -page 2057, column 1, paragraph 2; figures 3-8</p> <p>page 2061, column 1, paragraph 2 -page 2062, column 1, paragraph 1</p>	1-14
Y	<p>HOLDEN S ET AL: "Effects of ZD6474, an orally active inhibitor of VEGF receptor tyrosine kinase, in patients with solid tumors: Results from a phase I study."</p> <p>EUROPEAN JOURNAL OF CANCER, vol. 37, no. Supplement 6, October 2001 (2001-10), page S73</p> <p>XP002256123</p> <p>11th European Cancer Conference;Lisbon, Portugal; October 21-25, 2001, October, 2001</p> <p>ISSN: 0959-8049</p> <p>abstract</p>	1-14
Y	<p>HENNEQUIN LAURENT F ET AL: "Novel 4-anilinoquinazolines with C-7 basic side chains: Design and structure activity relationship of a series of potent, orally active, VEGF receptor tyrosine kinase inhibitors."</p> <p>JOURNAL OF MEDICINAL CHEMISTRY, vol. 45, no. 6, 14 March 2002 (2002-03-14), pages 1300-1312, XP002256124</p> <p>March 14, 2002</p> <p>ISSN: 0022-2623</p> <p>* see p. 1306-1307, ii.Inhbition of Endothelial Cell Proliferation ; vii.In Vivo Efficacy in Mouse ; Conclusions *</p> <p>abstract; figure 2; tables 4-6</p>	1-14

-/--

INTERNATIONAL SEARCH REPORT

PCT/GB 03/03390

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	"Activity of ZD6474, a vascular endothelial growth factor receptor tyrosine kinase inhibitor (VEGFR 'KDR!-TKI), in a model of ZD1839 ('Iressa') resistance" EUROPEAN JOURNAL OF CANCER, PERGAMON PRESS, OXFORD, GB, vol. 38, November 2002 (2002-11), pages S60-S61, XP004403633 ISSN: 0959-8049 abstract ---	1-14
P,Y	MANLEY PAUL W ET AL: "Therapies directed at vascular endothelial growth factor." EXPERT OPINION ON INVESTIGATIONAL DRUGS. ENGLAND DEC 2002, vol. 11, no. 12, December 2002 (2002-12), pages 1715-1736, XP002256125 ISSN: 1354-3784 abstract -----	1-14